

CYCLIC PERMUTATIONS IN KAZAMA-SUZUKI STRING MODELS

G. Aldazabal^{1,2,3}, I. Allekotte⁴, E. Andrés¹

and

C. Núñez^{2,5}

¹ *Centro Atómico Bariloche, 8400 S. C. de Bariloche,*

Comisión Nacional de Energía Atómica,

and Instituto Balseiro, Universidad Nacional de Cuyo, Argentina.

² *Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina*

³ *Universidad Autónoma, Madrid, Spain*

⁴ *Max-Planck-Institut für Physik komplexer Systeme*

Bayreuther Str. 40, Haus 16, 01187 Dresden, Germany

⁵ *Instituto de Astronomía y Física del Espacio*

C.C. 67, suc. 28, 1428 Buenos Aires

and Universidad de Buenos Aires, Argentina

(February 1, 2008)

Abstract

Moddings by cyclic permutation symmetries are performed in 4-dimensional strings, built up from $N = 2$ coset models of the type $CP_m = SU(m + 1)/SU(m) \times U(1)$. For some exemplifying cases, the massless chiral and antichiral states of E_6 are computed. The extent of the equivalence between different conformal invariant theories which possess equal chiral rings is analyzed.

I. INTRODUCTION

Algebraic constructions of 4 dimensional strings with $N = 1$ spacetime supersymmetry may be achieved by tensoring $N = 2$ superconformal field theories (see [1,2] and references therein). The $U(1)$ current of the $N = 2$ algebra provides the supersymmetry projector. Projection over odd integer charge states is required for modular invariance. The theory is anomaly free if the total internal central charge is $c_{int} = 9$.

Models constructed in this way possess a large set of invariances. Modding out by these symmetries, new modular invariant theories can be constructed. These orbifolds give rise to more acceptable models from the phenomenological point of view, in the sense that the number of chiral fermion generations is usually drastically reduced. A complete classification of these models could be useful for the discussion of interesting questions, such as mirror symmetry or topology change.

In particular, when two or more internal theories are equal, permutation symmetries are present. Actually, the first 3-generation (2,2) string model found by Gepner [3] is obtained when moddings by Z_3 phase and permutation symmetries are considered. A systematic study of cyclic permutation symmetries in Gepner's models was considered in [4,5] (see also [6]), where an interesting method to compute the massless spectrum of the orbifoldized theories was introduced.

Supersymmetric string vacua can also be constructed by orbifoldizing Landau-Ginzburg (LG) models [7–9]. A complete classification of LG superpotentials from which (2,2) string models can be constructed, including permutation symmetries of identical factors, was performed in [10]. When the LG superpotentials describe Gepner's models both methods yield the same number of E_6 generations. However in the case of LG models the chiral ring completely characterizes the theory, whereas the method introduced in [4,5] requires considering also non chiral states and allows to extract more information (for instance, the partition function of the permuted theory can be explicitly written).

In this note we study cyclic permutation symmetries in 4 dimensional string theories

built from more general $N=2$ superconformal coset models. In section II we recall the basics of CP_m cosets which are needed in our study. In section III we review the method to consider orbifolds of LG models with respect to abelian symmetries [11,12,10] and apply it to the computation of the number of E_6 generations of string models constructed with tensor products of cosets of the type $CP_m = SU(m+1)/SU(m) \times U(1)$. We then extend the method of references [4,5] to include non diagonal modular invariant couplings among left and right CP_m theories which do not admit a LG description. Even if the basic steps follow [4,5], the concrete computation demands a careful study of the component theory. In particular the construction of the cyclic permutation invariant states requires the knowledge of the exact weights and charges of the fields of the coset. A closed formula is known for the chiral (antichiral) fields but an each case computation is needed for the rest of the fields. In section IV we summarize the general method of references [4,5] and stress the distinct features of CP_m cosets. We illustrate the procedure of modding out by cyclic permutation symmetries of M theories (M prime) with two examples. The particular case $M = 2$ is analyzed in section V. Conclusions are presented in section VI. The explicit construction of the characters for $N = 2$ coset models is briefly reviewed in the Appendix.

II. $N = 2$ COSET MODELS

Here we consider the basic ingredients of CP_m coset models following the notation of [13]. For a more detailed discussion see for example [14].

We refer to the quotient theory $\frac{SU(m+1)_k \times SO(2m)_1}{SU(m)_{k+1} \times U(1)}$ as (m, k) . ω_i and $\hat{\omega}_i$ denote the fundamental weight vectors of $SU(m+1)$ and $SU(m)$, with i ranging from 0 to m and $m-1$ respectively. States of the $N = 2$ left superconformal algebra (SCA) are labelled by $|\Lambda, \lambda, \tilde{\Lambda}\rangle$, where Λ is a weight vector of $SU(m+1)$ at level k ($\Lambda = \sum_{i=1}^m m_i \omega_i$; $0 \leq \sum_{i=1}^m m_i \leq k$); $\tilde{\Lambda}$ is a $SO(2m)$ weight at level 1 (so it can only take the values 0, v , s , $\bar{s}$) and λ is a weight vector of $SU(m) \times U(1)$ at level $k+1$ obtained by decomposing $|\Lambda\rangle \otimes |\tilde{\Lambda}\rangle$ into irreducible representations of $SU(m) \times U(1)$.

A general field in the $N = 2$ quotient theory, denoted by $\Phi_{\Lambda\tilde{\Lambda}}^\lambda$, is obtained from the decomposition

$$G_\Lambda V_{\tilde{\Lambda}} = \Phi_{\Lambda\tilde{\Lambda}}^\lambda H_\lambda \quad (1)$$

where G_Λ , $V_{\tilde{\Lambda}}$, H_λ correspond to fields in the representations generated from $|\Lambda\rangle$, $|\tilde{\Lambda}\rangle$, $|\lambda\rangle$ respectively.

The conformal weight h and charge Q of $\Phi_{\Lambda\tilde{\Lambda}}^\lambda$ may in principle be obtained from this expression. In fact, by considering the $N = 2$ superconformal algebra associated to the Kac-Moody algebras $SU(m+1)$, $SO(2m)$ and $SU(m) \times U(1)$ it is easy to obtain the following equations

$$h = \frac{\Lambda(\Lambda + 2\rho_{m+1}) - \lambda(\lambda + 2\rho_m)}{2(k + m + 1)} + \frac{\tilde{\Lambda}^2}{2} + L \quad (2)$$

$$\begin{aligned} Q &= \sum_{l=1}^{2m} \tilde{\Lambda}^l - \frac{2}{k + m + 1} (\rho_{m+1} - \rho_m) \cdot \lambda + 2L' = \\ &= \sum_{l=1}^{2m} \tilde{\Lambda}^l - \frac{q}{k + m + 1} + 2L' \end{aligned} \quad (3)$$

where we have further decomposed λ into a $SU(m)$ weight $\hat{\lambda} = \sum_{i=1}^{m-1} \hat{n}_i \hat{\omega}_i$ and a $U(1)$ charge q (corresponding to the $U(1)$ of $SU(m) \times U(1)$) as:

$$\lambda = \hat{\lambda} + \frac{\omega_m}{m} q \quad (4)$$

ρ_{m+1} and ρ_m denote half the sum of the positive roots of $SU(m+1)$ and $SU(m)$ respectively. L, L' are integers which should be determined for each field of the theory, even for the primary ones [15]. The appearance of these integers relies in the fact that an irreducible representation of a Kac-Moody algebra $\hat{g}$ contains several representations of the original Lie algebra g . Every $\hat{g}$ -representation is generated by the application of Kac-Moody algebra generators $E_n^{\alpha_i}$, $H_n^{\alpha_i}$ to its highest weight. As $\hat{g}$ contains g as a subalgebra (generated by $E_0^{\alpha_i}$ and $H_0^{\alpha_i}$) then any representation of $\hat{g}$ can be decomposed into representations of g . The difference between the “ g -piece” of two weights of the $\hat{g}$ -representation is a root lattice vector of g . For some particular fields of the theory it is possible to write down a closed

formula for h and Q . In fact, for the chiral fields corresponding to $\Lambda = \lambda$ and $\tilde{\Lambda} = 0$, $h = Q/2$ is obtained from the above equations with $L = L' = 0$ [15] (and similarly for the antichiral fields with $\Lambda = \lambda^*$, where the asterisk means conjugate representation).

In the study of permutation symmetries the exact weights and charges of all the primary fields are necessary, i.e. L and L' must be determined. They are computed in some examples in section IV by explicit construction of the corresponding characters, following [16]. This is essentially done by decomposing the characters χ_Λ^G , $\chi_{\tilde{\Lambda}}^{SO(d)}$ into the characters χ_λ^H of $SU(m) \times U(1)$. The character of the coset conformal theory is defined through

$$\chi_\Lambda^G \chi_{\tilde{\Lambda}}^{SO(d)} = \sum_{\lambda} \chi_{\Lambda, \lambda, \tilde{\Lambda}}^{N=2} \chi_\lambda^H. \quad (5)$$

When counting primary states proper external automorphisms of the affine Kac-Moody algebra must be taken into account. Under the automorphism σ a state $|\Lambda, \hat{\lambda}, q, \tilde{\Lambda}\rangle$ changes to [17–19] $|\sigma(\Lambda), \sigma(\hat{\lambda}), \sigma(q), \sigma(\tilde{\Lambda})\rangle$, where

$$\begin{aligned} \sigma(\Lambda) &= (k - \sum_{i=1}^m n_i) \omega_1 + \sum_{i=2}^m n_{i-1} \omega_i \\ \sigma(\hat{\lambda}) &= (k + 1 - \sum_{i=1}^{m-1} n_i) \omega_1 + \sum_{i=2}^{m-1} n_{i-1} \omega_i \\ \sigma(q) &= q + k + m + 1 \\ \sigma(\tilde{\Lambda} = (0), (v), (s), (\bar{s})) &= ((v), (0), (\bar{s}), (s)) \end{aligned} \quad (6)$$

When applying σ simultaneously to left and right movers the resulting states must be identified so that the fields in the theory form a unitary representation of the modular group. Further identifications are required, depending on the symmetry properties of the modular invariants $\mathcal{N}$, $\mathcal{M}$ defined below [13]. As the CP_m models do not possess automorphism fixed points, this field identification poses no further problem (for a discussion of fixed points in coset models see [20,21]).

Finally, the full partition function of the CP_m $N = 2$ superconformal theory is

$$Z = \sum_{\Lambda, \lambda, \tilde{\Lambda}, \bar{\lambda}, \bar{\tilde{\Lambda}}} \chi_{\Lambda, \lambda, \tilde{\Lambda}}^{N=2} \mathcal{N}_{\Lambda, \tilde{\Lambda}} \mathcal{M}_{\lambda, \bar{\lambda}} \chi_{\tilde{\Lambda}, \bar{\lambda}, \bar{\tilde{\Lambda}}}^{N=2*} \quad (7)$$

In the above equation the sum extends over states (Λ, λ) and $(\bar{\Lambda}, \bar{\lambda})$ satisfying the condition $C(\Lambda, \lambda)$ of [17], which for the Neveu-Schwarz (NS) sector corresponds to $\Lambda - \lambda \in M$, the root lattice of $SU(m+1)$. $\mathcal{N}$ and $\mathcal{M}$ denote modular invariants for $SU(m+1)$ and $SU(m) \times U(1)$ respectively. Throughout the paper, barred (unbarred) quantities denote right (left) movers.

To make a string theory out of these models the Gepner construction has to be followed [1]: each sector (left and right moving) will be a product of spacetime bosons and fermions, times the product of r internal $N = 2$ coset fields, such that $c_{int} = 9$. To obtain $N = 1$ spacetime supersymmetry, a projection over states with odd integer $U(1)$ charge Q must be performed.

It is useful to denote each state in the full theory by a vector

$$V = (\tilde{\Lambda}_0; q_1, \dots, q_r; \tilde{\Lambda}_1, \dots, \tilde{\Lambda}_r) \quad (8)$$

where $\tilde{\Lambda}_0$ is a $SO(2)$ weight. We will also need the definition of a scalar product

$$V \cdot V' = \sum_{i=0}^r \tilde{\Lambda}_i \tilde{\Lambda}'_i - \sum_{i=1}^r \frac{q_i q'_i}{2\eta_i(k_i + m_i + 1)} \quad (9)$$

and the vectors

$$\begin{aligned} \beta_0 &= (\bar{s}; \eta_1, \dots, \eta_r; s_1, \dots, s_r) \\ \beta_i &= (v; 0, \dots, 0; 0, \dots, v, \dots, 0) \quad (v \text{ in the } i\text{-th. position}) \end{aligned} \quad (10)$$

Here $\eta_i = \frac{1}{2}m_i(m_i + 1)$.

The supersymmetry projection and aligned boundary conditions, Ramond-Ramond (R-R) and NS-NS, are now accomplished by

$$Q(\bar{V}) = 2\beta_0 \cdot \bar{V} = \text{odd integer} \quad (11)$$

$$2\beta_i \cdot \bar{V} = \text{even integer} \quad (12)$$

For NS states condition (11) amounts to integer internal charge Q_{int} . To maintain modular invariance, twisted sectors must be included. These are given by the condition

$$V = \bar{V} + s\beta_0 + \sum_{i=1}^r n_i \beta_i \quad (13)$$

where s and n_i are integers. Of course a given state is allowed whenever both invariants $\mathcal{N}$ and $\mathcal{M}$ are nonvanishing and its multiplicity is given by the product of the modular coefficients.

The heterotic construction is implemented by replacing left spacetime fermions by internal free bosons with central charge $c = 24$ to cancel the bosonic anomaly. Modular invariance of the theory follows from the isomorphism between representations of $SO(2)$ and the new gauge group $E_8 \times SO(10)$ under the modular group. With the supersymmetry projection the gauge group is enlarged from $E_8 \times SO(10)$ to $E_8 \times E_6$.

As is well known [1,22,6,13], CP_m coset models have a discrete Z_{k+m+1} symmetry. New models can be built from the original one modding by this symmetry; corresponding string compactifications may have a smaller number of generations.

These moddings are characterized by a vector

$$\Gamma = (0; \gamma_1 \eta_1, \dots, \gamma_r \eta_r; 0, \dots, 0) \quad (14)$$

where γ_i are integers satisfying

$$2\beta_0 \cdot \Gamma = - \sum_{i=1}^r \frac{\gamma_i \eta_i}{k_i + m_i + 1} = integer \quad (15)$$

States in the twisted sectors now verify

$$V = \bar{V} + s\beta_0 + \sum_{i=1}^r n_i \beta_i + 2x\Gamma \quad (16)$$

and the generalized GSO projection over states with integer Q_{int} is now given by the conditions (11) and (12) and

$$- \Gamma \cdot (2\bar{V} + 2x\Gamma) = integer \quad (17)$$

(see [6]). The modular invariant partition function is given by

$$Z = \frac{1}{DM_i} \sum_{n_i, m_i, x, y, t, s} Z(s, n_i, x, t, m_i, y) \quad (18)$$

with

$$\begin{aligned}
Z(s, n_i, x, t, m_i, y) = \\
= \sum_V (-1)^{s+t} e^{-2i\pi V(t\beta_0 + \sum_i m_i \beta_i)} e^{-2i\pi y \Gamma(2V + 2x\Gamma + 2s\beta_0)} \chi_V \chi_{V+s\beta_0 + \sum_i n_i \beta_i + 2x\Gamma}^*
\end{aligned} \tag{19}$$

Summations over x and y range from 0 to $M - 1$, M being the least integer such that $M\gamma_i \in Z$ for all γ_i . The sum over y implements condition (17). Again, only states allowed by the invariants $\mathcal{M}$ and $\mathcal{N}$ are coupled.

III. CYCLIC PERMUTATIONS IN LANDAU GINZBURG MODELS

When considering CP_m coset models with diagonal modular invariants, the computation of chiral fermion generations can be simplified by exploiting the known relation to $N = 2$ LG models. We refer the reader to references [7–9] for the basics of LG description of superconformal $N = 2$ theories and here we concentrate on the calculation of the number of chiral generations in the corresponding superstring vacua. The LG action in superspace is

$$S = \left(\int d^2 z d^4 \theta K(\phi_i, \bar{\phi}_i) \right) + \left(\int d^2 z d^2 \theta W(\phi_i) + c.c. \right) \tag{20}$$

The superpotential W is holomorphic in the chiral fields ϕ_i and, in the fixed point of the renormalization group trajectories, defines a superconformal theory with quasihomogeneous potential

$$W(\lambda^{n_i} \phi_i) = \lambda^D W(\phi_i) \tag{21}$$

In order to describe a $N = 2$ string vacuum from a LG theory, the GSO conditions on integer charge states must be imposed. This corresponds to considering the orbifold by the discrete group transforming the fields as

$$\Phi_i \rightarrow e^{2\pi i n_i / D} \Phi_i \tag{22}$$

where n_i are the weights and D is the order of the superpotential.

The number of fermionic generations contained in this theory can be computed with the following formula derived in [11,12],

$$N_{gen} = -\frac{1}{2D} \sum_{p,s=0}^{D-1} (-)^{(\frac{c}{3}-r)(p+s+ps)} \prod_{pQ_i, sQ_i \in Z} \frac{n_i - D}{n_i} \quad (23)$$

where the product is taken over all n_i such that $pQ_i \in Z$ and $sQ_i \in Z$ simultaneously.

In general, LG models possess a larger discrete symmetry group than that generated by (22). They have been classified in reference [10] where string vacua as orbifolds by discrete abelian symmetries were constructed. In a base in which all the elements of the symmetry group act diagonally on the fields it is possible to characterize each element g of a cyclic group G by the order M of g , and by an integer r -component vector $\gamma = (\gamma_1, \dots, \gamma_r)$

$$g : \phi_i \rightarrow e^{2\pi i \Theta_i^g} \phi_i = e^{2\pi i \gamma_i / M} \phi_i \quad (24)$$

When $\det(g) = 1$ for all $g \in G$, the orbifold of the LG model is (2,2) supersymmetric and still has an interpretation as a Calabi-Yau or an orbifold of a Calabi-Yau manifold compactification.

The number of generations of a theory orbifoldized by n cyclic groups characterized by $\gamma^{(1)}, \dots, \gamma^{(n)}$ is [12]

$$N_{gen} = -\frac{1}{2 \prod_{i=0}^n M_i} \sum_{p_i, s_i=0}^{M_i-1} \prod_{\substack{j/ \\ \sum_i p_i \gamma_j^{(i)} / M_i \in Z \\ \sum_i s_i \gamma_j^{(i)} / M_i \in Z}} \left(1 - \frac{D}{n_j}\right) \quad (25)$$

where $M_0 = D, p_0 = p, s_0 = s$ and $g^{(0)} = (n_1, \dots, n_r)$. This expression does not include discrete torsion and holds for $(\frac{c}{3} - r)$ even. A trivial field may be added whenever this condition is not satisfied, namely $W' = W + \phi_t^2$.

In order to orbifoldize by discrete symmetries using this formalism, notice that if a generic potential W has a cyclic permutation symmetry of M fields (M prime), namely

$$W(\phi_0, \dots, \phi_{M-1}) = W(\phi_{M-1}, \phi_0, \dots, \phi_{M-2}) \quad (26)$$

it is possible to perform the following change of variables

$$\psi_j = \frac{1}{\sqrt{N}} \sum_{k=0}^{M-1} e^{-2\pi i k j / M} \phi_k \quad (27)$$

which trades a permutation symmetry of the fields ϕ_k into a phase symmetry of ψ_k (since $\phi_k \rightarrow \phi_{k+1}$ implies $\psi_j \rightarrow e^{2\pi i j/M} \psi_j$). Notice that ϕ_k and ψ_k have the same weight, and thus the orbifold by β_0 is not modified by the change of variables. The number of generations of the theory can thus be computed with eqn. (25) choosing $\gamma_0 = (n, \dots, n, n_M, \dots, n_r)$ and $\gamma^{(1)} = (0, 1, 2, \dots, M-1, 0, \dots, 0)$ of order M . The M prime condition implies

$$\det(g) = e^{2\pi i \sum_{j=0}^{M-1} \frac{j}{M}} = 1 \quad (28)$$

Notice that it is not necessary to know the superpotential explicitly in order to compute N_{gen} in the orbifoldized theory. Knowledge of the weights and the order of W suffices.

By applying this procedure to all $N = 2$ Gepner models we reobtained the results of reference [5]. The same calculation was performed for CP_m Kazama-Suzuki models that can be described in terms of a LG superpotential. These are the diagonal $(m, k)_{AA}$ (the subindices denote the $\mathcal{N}$ and $\mathcal{M}$ modular invariants respectively) which have [23,9]

$$W = \sum_{j_1+2j_2+\dots+mj_m=k+m+1} A_{n_1\dots n_m} \phi_1^{j_1} \dots \phi_m^{j_m} \quad (29)$$

where the $A_{n_1\dots n_m}$ coefficients are such that each ϕ_i has weight j .

The results are listed in Table 1 where the corresponding superpotentials are denoted by D, $(n_1, \dots, n_r)$ and the cyclic permutations considered can be deduced from the vectors γ . We have not included models known to be equivalent at the chiral level because they clearly yield the same results. Notice that eqn.(25) allows to perform the sum over the p_i 's separately in each sector $(s_0, \dots, s_n)$. Thus the contribution to N_{gen} can be obtained sector by sector. However, for non-LG models another method has to be applied. We describe it in the following sections, where we also emphasize other by-products of this formulation.

TABLE I

Number of generations in CP_m models orbifoldized by cyclic permutation symmetries

Model	$D, (n_1, \dots, n_r)$	M, γ	N_{gen}
$(2, 3)_{AA} \times (2, 3)_{AA} \times (2, 3)_{AA}$	$6, (1, 2, 1, 2, 1, 2)$	$3, (0, 0, 1, 1, 2, 2)$	36
$(2, 3)_{AA} \times (2, 3)_{AA} \times (2, 3)_{AA}$	$6, (1, 2, 1, 2, 1, 2)$	$2, (0, 0, 1, 1, 0, 0)$	54
$(4, 5)_{AA} \times 1_A \times 1_A \times 1_A$	$30, (3, 6, 9, 12, 10, 10, 10)$	$3, (0, 0, 0, 0, 0, 1, 2)$	0
$(3, 4)_{AA} \times 1_A \times 1_A \times 1_A \times 2_A$	$24, (3, 6, 9, 8, 8, 8, 6)$	$3, (0, 0, 0, 0, 1, 2, 0)$	0
$(3, 3)_{AA} \times 1_A \times 1_A \times 1_A \times 5_A$	$21, (3, 6, 9, 7, 7, 7, 3)$	$3, (0, 0, 0, 0, 1, 2, 0)$	0
$(3, 4)_{AA} \times 2_A \times 2_A \times 2_A$	$8, (1, 2, 3, 2, 2, 2)$	$3, (0, 0, 0, 0, 1, 2)$	40
$(3, 4)_{AA} \times 6_D \times 6_D$	$8, (1, 2, 3, 2, 3, 2, 3)$	$2, (0, 0, 0, 0, 0, 1, 1)$	40
$(3, 3)_{AA} \times 12_D \times 12_D$	$7, (1, 2, 3, 1, 3, 1, 3)$	$2, (0, 0, 0, 0, 0, 1, 1)$	56
$(2, 4)_{AA} \times (2, 4)_{AA} \times 5_A$	$7, (1, 2, 1, 2, 1)$	$2, (0, 0, 1, 1, 0)$	51
$(2, 6)_{AA} \times (2, 6)_{AA} \times 1_A$	$9, (1, 2, 1, 2, 3)$	$2, (0, 0, 1, 1, 0)$	42
$(2, 6)_{AA} \times 1_A \times 1_A \times 1_A \times 1_A \times 1_A$	$9, (1, 2, 3, 3, 3, 3, 3)$	$3, (0, 0, 0, 1, 2, 0, 0)$	60
$(2, 6)_{AA} \times 1_A \times 1_A \times 1_A \times 1_A \times 1_A$	$9, (1, 2, 3, 3, 3, 3, 3)$	$5, (0, 0, 0, 1, 2, 3, 4)$	12
$(2, 6)_{AA} \times 1_A \times 1_A \times 1_A \times 1_A \times 1_A$	$9, (1, 2, 3, 3, 3, 3, 3)$	$2, (0, 0, 0, 1, 0, 1, 0)$	48
$(2, 6)_{AA} \times 1_A \times 1_A \times 1_A \times 4_A$	$18, (2, 4, 6, 6, 6, 3)$	$3, (0, 0, 0, 1, 2, 0)$	60
$(2, 4)_{AA} \times 1_A \times 1_A \times 1_A \times 12_A$	$42, (6, 12, 14, 14, 14, 3)$	$3, (0, 0, 0, 1, 2, 0)$	0
$(2, 6)_{AA} \times 1_A \times 1_A \times 2_A \times 2_A$	$36, (4, 8, 6, 15, 6, 15)$	$2, (0, 0, 0, 0, 1, 1)$	0
$(2, 4)_{AA} \times 26_D \times 26_D$	$28, (4, 8, 2, 13, 2, 13)$	$2, (0, 0, 0, 0, 1, 1)$	102

IV. CYCLIC PERMUTATIONS IN CP_m COSETS

Cyclic permutation symmetries in Gepner models were systematically considered in [4,5]. In this section we extend the method developed in these references to more general CP_m coset theories.

Assume that there are M identical CP_m blocks in the internal sector of the 4-D string. We consider a prime number of identical theories, but the analysis can be easily extended to any M . Let us introduce a formal projection operator P over identical states of the $N = 2$ superconformal algebra (not necessarily primary states), such that P acting on a tensor product of states produces a vanishing result unless all states have equal charges and weights. After dividing by this permutation symmetry the following “character” can be defined

$$\begin{aligned}\chi_{invar}(\tau) &= (P + (1 - P)/M)\chi(\tau) = \frac{\chi_{original}}{M} + \frac{M - 1}{M}P\chi_{original} = \\ &= \frac{\chi^M(\tau)}{M} + \frac{M - 1}{M}\chi(M\tau)\end{aligned}\tag{30}$$

$P\chi$ formally indicates that the traces must be computed by simultaneously considering the same state in all blocks, such that $P\chi(\tau) = \chi(M\tau)$. Each of these states is counted once. The term $\frac{(1-P)}{M}$ corresponds to the case when at least one state in a block is different from the others. Because this state could belong to any of the M blocks we must divide by M in order to obtain just one full symmetric state. $\chi_{invar}(\tau)$ does not transform properly under modular transformations and twisted sectors must be added. The full modular invariant partition function of the projected theory may finally be written in terms of the character of just one of the (identical) component theories

$$Z_{new}(\tau, \bar{\tau}) = \frac{Z^M(\tau, \bar{\tau})}{M} + \frac{M - 1}{M}Z(M\tau, M\bar{\tau}) + \frac{M - 1}{M} \sum_{n=0}^{M-1} Z\left(\frac{\tau + n}{M}, \frac{\bar{\tau} + n}{M}\right)\tag{31}$$

Therefore, if fields of the original CP_m theory are known, (31) allows to compute the spectrum in the modded theory. We shall refer to the states coming from the last term as twisted states. An analysis of this term shows that it can be interpreted as the partition function

of a new $N = 2$ superconformal field theory with central charge $\hat{c} = Mc$ (c is the central charge of each one of the identical theories) and Virasoro generators given in terms of those of the original theory by

$$\hat{L}_m = \frac{L_{mM}}{M} + \frac{c(M^2 - 1)}{24M} \delta_{0,m} \quad (32)$$

$$\hat{G}_r^\pm = \frac{1}{\sqrt{M}} G_{rM}^\pm \quad (33)$$

$$\hat{J}_m = J_{mM} \quad (34)$$

Therefore the weights and charges of the (twisted) primary states of the new theory are obtained from the original ones as

$$h_{new} = \frac{h + m}{M} + \frac{c(M^2 - 1)}{24M} \quad (35)$$

$$Q_{new} = Q \quad (36)$$

where m is the level of the descendant field. Similar expressions are valid for the right movers.

The sum over n in (31) imposes the constraint $h + m - \bar{h} - \bar{m} = 0 \pmod{M}$.

In order to build up the partition function for the 4-D string the spacetime sector and the other $r - M$ CP_m blocks must be included. Orbifolds by β_0 and β_i must then be performed. Orbifolds by commuting phase symmetries could also be considered.

Following the reasoning of ref. [5], the Z_{new} corresponding to (31) is constructed in terms of the partition function written in (18). Modular transformations allow to build the twisted sector. The conclusion is that (11), (12) and (17) are still valid in the twisted sector if we define the new $1 + 2(r - M + 1)$ component vector

$$\begin{aligned} V &= (\tilde{\Lambda}_0; q_1, q_{M+1}, \dots, q_r; \tilde{\Lambda}_1, \tilde{\Lambda}_{M+1}, \dots, \tilde{\Lambda}_r) \\ \beta_0 &= (\bar{s}; M\eta_1, \eta_{M+1}, \dots, \eta_r; s_1, s_{M+1}, \dots, s_r) \\ \beta_i &= (v; 0, \dots, 0; 0, \dots, v, \dots, 0) \end{aligned} \quad (37)$$

and similarly for the phase vector

$$\Gamma = (0; M\gamma_1\eta_1, \gamma_{M+1}\eta_{M+1}, \dots, \gamma_r\eta_r; 0, \dots, 0) \quad (38)$$

where the M original identical theories have been replaced by just one in the first block.

Eqn. (16) now means that NS left states may couple to NS right states if

$$q_1 - \bar{q}_1 = m_1(m_1 + 1)M(s + x\gamma_1) \quad \text{mod} \quad m_1(m_1 + 1)M(k_1 + m_1 + 1) \quad (39)$$

$$q_i - \bar{q}_i = m_i(m_i + 1)(s + x\gamma_i) \quad \text{mod} \quad m_i(m_i + 1)(k_i + m_i + 1) \quad (40)$$

for $i = M + 1, \dots, r$.

The issue of field identifications in the modded theory deserves a comment. In fact, in the new theory, expressed in terms of just one original block, the orbit under σ has been enlarged up to $M(m_1 + 1)m_1$. An indication of it is that if we compute the conformal weight h in (35) by just using the σ transformed state (7) the weights in the twisted sector (16) will be $h_{new} + \text{integer}$, only after applying $Mm_1(m_1 + 1)$ times the transformation σ [19].

Massless chiral-chiral (chiral-antichiral) matter (antimatter) fields satisfy $h_{int} = Q_{int}/2$ and $\bar{h}_{int} = \bar{Q}_{int}/2$ ($\bar{h}_{int} = -\bar{Q}_{int}/2$). The chiral (antichiral) states of the tensor product of models are obtained as products of chiral (antichiral) states of each coset. Similarly for the new theory, chiral states have $h_{new} = Q_{new}/2 = Q/2$.

We will apply the method discussed above to mod out by cyclic permutations of $M = 3$ blocks in the $(2, 1)_{AA}^4$ theory ($c_{int} = 6$) and in the $(2, 2)_{AA}^3 3_A$ theory ($c_{int} = 9$) and compute the massless chiral spectrum in both cases (**56** and **27** and **27** of E_6 respectively). We indicate the permuted theories in brackets, i.e. $\{(2, 1)_{AA}^3\}(2, 1)_{AA}$ and $\{(2, 2)_{AA}^3\}3_A$ respectively.

According to notations of section II, in a general $(2, k)$ theory

$$\Lambda = m_1 w_1 + m_2 w_2 \quad (41)$$

$$\lambda = n \hat{w}_1 + \frac{q}{2} w_2 \quad (42)$$

such that a state in the coset may be written as $(m_1, m_2)(n, q)s$ (with $s = 0, 2, 1, -1$ corresponding to $0, v, s, \bar{s}$). The constraints $C(\Lambda, \lambda)$ read in the NS sector

$$m_1 + 2m_2 - q = 0 \quad \text{mod} \quad 3$$

$$4m_1 + 2m_2 - q - 3n = 0 \quad \text{mod} \quad 6$$

(with $m_1 + m_2 \leq k$ and $q \equiv q + m(m+1)(k+m+1)$) and the weights and charges are

$$h = \frac{1}{12(k+m+1)}[4m_1^2 + 4m_2^2 + 4m_1m_2 + 12m_1 + 12m_2 - 3n(n+2) - q^2] + \frac{s^2}{8} + L \quad (43)$$

$$Q = \frac{-q}{k+m+1} + \frac{s}{2} + 2L'. \quad (44)$$

Consider the $(2,1)_{AA}$ coset. For this case, when $M = 3$, the above conditions lead to six highest weight states, namely

state #	(m_1, m_2)	n	q	s	h	Q	h_{new}
1	(0, 0)	0	0	0	0	0	$\frac{1}{6}$
2	(0, 1)	0	2	0	$\frac{1}{4}$	$\frac{-1}{2}$	$\frac{1}{4}$
3	(1, 0)	0	-2	0	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$
4	(1, 0)	1	1	0	$\frac{1}{8}$	$\frac{-1}{4}$	$\frac{5}{24}$
5	(0, 1)	1	-1	0	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{5}{24}$
6	(1, 0)	0	4	2	$\frac{1}{2} + L$	$2L'$	$\frac{1+L}{3}$

and their corresponding σ transformed states (7). Note that the integers L, L' entering the definitions (43) and (44) are unknown only for the sixth field, as this is the only one which is neither chiral nor antichiral. In order to exactly evaluate them we look at the corresponding character (see appendix) expressed as an expansion in powers of $x = e^{2i\pi\tau}$ and $y = e^{2i\pi z}$. (Note that only the first non vanishing power of x and y respectively are necessary to determine these integers).

The first terms of the character for this state are

$$\chi_{(0,4)}^{(1,0)} = x^{1/2}y^0[1 + (y + y^{-1})x^{1/2} + 2x + (y + y^{-1})x^{3/2} + 4x^2 \dots] \quad (45)$$

Therefore, $h = 1/2$ and $Q = 0$. The next power of x is increased in $1/2$. This is a reflection of the general fact that the original $N = 2$ superconformal algebra may be split into subalgebras

built up by the application of an even or an odd number of currents $G_{-1/2}^{\pm}$ to the original primary state (see [1]). The states obtained from the primary by one of these subalgebras are labeled with $s = 0$, and those obtained by the other, with $s = 2$, respectively. Since the σ transformation interchanges $s = 0$ and $s = 2$, it is possible that a primary state has $s = 2$ as above (notice that the state $((1,0)(0,4)2)$ is identified with $((0,1)(2,0)0)$). As a check of our computations it can be shown that the other fields in the σ orbit produce the same character.

Notice that in the case at hand the integers L and L' could have been guessed by claiming the duality relation (valid after field identifications) $(m, k) = (k, m)$ [2] so that $(2, 1)_{AA}$ corresponds to the well known minimal model 2_A . We have explicitly checked this relation by comparing the characters of the two models up to several powers of x and y .

We now turn our attention to cyclic permutations. We are interested in computing the number of chiral-chiral and chiral-antichiral states in the projected theory. From (35) and (36) we find that chiral (and antichiral) states in the twisted sector come from states in the σ -orbits of

<i>state #</i>	$((m_1, m_2)(n, q)s)$	h	Q	h_{new}
2	$((0, 1)(0, 2)0)$	$\frac{1}{4}$	$\frac{-1}{2}$	$\frac{1}{4}$
3	$((1, 0)(0, -2)0)$	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$
7	$G_{-1/2}^+((1, 0)(0, 4)2)$	1	1	$\frac{1}{2}$
8	$G_{-1/2}^-((1, 0)(0, 4)2)$	1	-1	$\frac{1}{2}$
9	$G_{-1/2}^-((0, 1)(1, -1)0)$	$\frac{5}{8}$	$\frac{-3}{4}$	$\frac{3}{8}$
10	$G_{-1/2}^+((1, 0)(1, 1)0)$	$\frac{5}{8}$	$\frac{3}{4}$	$\frac{3}{8}$

Attaching the chiral states of the fourth $(2, 1)_{AA}$ theory (i.e. the theory which is not permuted) and coupling left and right movers according to (11), (12) and (17) we obtain

the following chiral-chiral states in the twisted sector.

$$\begin{array}{rclcl}
\{7\} & 1 & - & \{7\} & 1 \\
\{10\} & 5 & - & \{10\} & 5 \\
\{3\} & 3 & - & \{3\} & 3 \\
\{7\} & 1 & - & \sigma^6(\{3\}) & \sigma^2(3) \\
\{10\} & 5 & - & \sigma^9(\{10\}) & \sigma^3(5) \\
\{3\} & 3 & - & \sigma^{12}(\{7\}) & \sigma^4(1)
\end{array}$$

where the numbers denote the state # in the tables above.

By adding the states in the untwisted sector we finally obtain a total of 20 states. This coincides with the result found in [5] for the $\{2_A\}^3 2_A$ model, as expected.

Let us now discuss the second example $\{(2, 2)_{AA}^3\} 3_A$. The $(2, 2)_{AA}$ coset and the 8_D minimal model are known to have the same spectrum [14,24], so their Poincaré polynomials coincide. However the full conformal theories are different in the sense that the 8_D theory has 25 primary states of the Virasoro algebra while $(2, 2)_{AA}$ has 20. For example the primaries with (l, q, s) and $(l, -q, s)$ equal to $(6, 4, 0)$, $(6, 2, 0)$, $(6, 0, 0)$, $(8, 4, 0)$ and $(8, 0, 0)$ are states of 8_D but there do not exist corresponding ones with the same weight and charge in $(2, 2)_{AA}$. Note that the states with $l = 4$ have multiplicity 2 due to the modular invariant. Moreover, the characters of some of the chiral primary fields in 8_D and $(2, 2)_{AA}$ (with the same conformal weights) turn out to be different. For example the character of the identity in 8_D is

$$\begin{aligned}
\chi_{0,0} = & 1 + x + 3x^2 + 6x^3 + x^4(13 + y^2 + y^{-2}) + x^{3/2}(y + y^{-1}) + \\
& + 2x^{5/2}(y + y^{-1}) + 5x^{7/2}(y + y^{-1}) + \dots
\end{aligned} \tag{46}$$

while in $(2, 2)_{AA}$ it is

$$\begin{aligned}
\chi_{0,0}^{0,0} = & 1 + x + 4x^2 + 8x^3 + x^4[19 + 2(y^{-2} + y^2)] + x^{3/2}(y^{-1} + y) + \\
& + 3x^{5/2}(y^{-1} + y) + 7x^{7/2}(y^{-1} + y) + \dots
\end{aligned} \tag{47}$$

indicating that both fields transform differently under the conformal algebra. States that are neither chiral nor antichiral may contribute to form the twisted chiral states in the new

theory (see (35), (36)) and because these states are not the same for both theories, the number of states in the twisted sectors may be expected to differ. However, this is not the case in the model we are discussing.

The new chiral (antichiral) states in the twisted sector are

<i>state</i>	<i>h</i>	<i>Q</i>	<i>h_{new}</i>
$(2, 0)(0, \pm 4)0$	$\frac{2}{5}$	$\mp \frac{4}{5}$	$\frac{2}{5}$
$G_{-1/2}^{\pm}((0, 1)(1, 5)2)$	$\frac{7}{10}$	± 1	$\frac{1}{2}$

Attaching the chiral (antichiral) states of 3_A such that the total weight and charge add up to $1/2$ and 1 (-1) we obtain 4 states in the twisted sector of the **27** and also of the $\overline{\mathbf{27}}$. The same number of twisted states was obtained in [5] for the $\{8_D^3\}3_A$ model. This coincidence was not expected a priori since the length of the σ orbits of both theories are different. However since the left and right movers have to couple in the $(2, 2)_{AA}$ model according to the $\mathcal{N}_{\Lambda, \bar{\Lambda}}$ and $\mathcal{M}_{\lambda, \bar{\lambda}}$ modular invariants, only 3 states in the σ chain potentially contribute to the twisted sector similarly as in the 8_D case.

In the example we are discussing the equivalent results for both theories can be explained by noticing that, even if the fields of both models transform differently under the action of the conformal generators, the partition functions turn out to be equal. In fact we have explicitly checked that some characters of the $(2, 2)_{AA}$ are sum of characters of the 8_D , and they are so in a way that the partition functions of both models coincide. The remaining characters of the 8_D and the $(2, 2)_{AA}$ models are equal one to one (so the corresponding states transform in the same way under the conformal group).

V. Z_2 PERMUTATIONS

As discussed in [5] care needs to be taken when considering permutations of one pair of blocks. Since states with $s = 2$ ($\bar{s} = 2$) can be obtained by acting with $G_{-1/2}^{\pm}$ ($\bar{G}_{-1/2}^{\pm}$) on the corresponding states with $s = 0$ ($\bar{s} = 0$), permuting a pair of states both having one of these fermionic operators introduces a minus sign which amounts to the cancellation of the

symmetrized state in the spectrum. Of course this sign will not appear if an even number of permutations of two theories is considered, the cancellation does not take place and (31) directly applies.

The twisted sectors, as shown in [5], have to be constructed in the R sector of the original theory in the case of Z_2 permutations.

As an example we study $M = 2$ permutations in $(2, 1)^4$. Consider first two permutations of two blocks, i.e. $\{(2, 1)^2\}\{(2, 1)^2\}$. By applying the spectral flow shift β_0 to the NS states we obtain the following set of R states

<i>block state</i>	<i>h</i>	<i>h_{new}</i>	<i>Q</i>
(0, 0)(0, 3)1	1/16	1/8	1/4
(0, 1)(0, 5)1	1/16	1/8	-1/4
(1, 0)(0, 1)1	9/16	3/8	3/4
(1, 0)(1, 4)1	1/16	1/8	0
(0, 1)(1, 2)1	5/16	1/4	1/2
(1, 0)(0, 7) - 1	9/16	3/8	1/4

Coupling left and right movers according to (39) we obtain the following chiral-chiral states in the twisted sector

$$\begin{aligned}
& \{(0, 0)(0, 3)1\} \{(1, 0)(0, 1)1\} - \{(0, 0)(0, 3)1\} \{(1, 0)(0, 1)1\} \\
& \{(0, 0)(0, 3)1\} \{(1, 0)(0, 1)1\} - \sigma^6 \{(0, 0)(0, 3)1\} \sigma^6 \{(1, 0)(0, 1)1\} \\
& \{(1, 0)(0, 1)1\} \{(0, 0)(0, 3)1\} - \{(1, 0)(0, 1)1\} \{(0, 0)(0, 3)1\} \\
& \{(1, 0)(0, 1)1\} \{(0, 0)(0, 3)1\} - \sigma^6 \{(1, 0)(0, 1)1\} \sigma^6 \{(0, 0)(0, 3)1\} \\
& \{(0, 1)(1, 2)1\} \{(0, 1)(1, 2)1\} - \{(0, 1)(1, 2)1\} \{(0, 1)(1, 2)1\} \\
& \{(0, 1)(1, 2)1\} \{(0, 1)(1, 2)1\} - \sigma^3 \{(0, 1)(1, 2)1\} \sigma^3 \{(0, 1)(1, 2)1\} \\
& \{(0, 1)(1, 2)1\} \{(0, 1)(1, 2)1\} - \sigma^6 \{(0, 1)(1, 2)1\} \sigma^6 \{(0, 1)(1, 2)1\} \\
& \{(0, 1)(1, 2)1\} \{(0, 1)(1, 2)1\} - \sigma^9 \{(0, 1)(1, 2)1\} \sigma^9 \{(0, 1)(1, 2)1\}
\end{aligned}$$

I.e., a total of eight twisted states. Since this model has four invariant states, applying (31) leads to a number of 20 chiral-chiral states as is consistent with a $c_{int} = 6$ model.

Let us now consider the case $\{(2,1)^2\} \ (2,1)^2$, i.e. permutations of only one pair of blocks. We have to combine the R states of the “new” theory with the NS states of the two unpermuted theories. Again we obtain eight states in the twisted sector. They are

$$\begin{aligned}
& \{(0,0)(0,3)1\}((0,1)(1,-1)0)((1,0)(0,-2)0) - \{(0,0)(0,3)1\}((0,1)(1,-1)0)((1,0)(0,-2)0) \\
& \{(0,0)(0,3)1\}((1,0)(0,-2)0)((0,1)(1,-1)0) - \{(0,0)(0,3)1\}((1,0)(0,-2)0)((0,1)(1,-1)0) \\
& \{(1,0)(0,1)1\}((0,1)(1,-1)0)((0,0)(0,0)0) - \{(1,0)(0,1)1\}((0,1)(1,-1)0)((0,0)(0,0)0) \\
& \{(1,0)(0,1)1\}((0,0)(0,0)0)((0,1)(1,-1)0) - \{(1,0)(0,1)1\}((0,0)(0,0)0)((0,1)(1,-1)0) \\
& \{(0,1)(1,2)1\}((0,1)(1,-1)0)((0,1)(1,-1)0) - \{(0,1)(1,2)1\}((0,1)(1,-1)0)((0,1)(1,-1)0) \\
& \{(0,1)(1,2)1\}((0,0)(0,0)0)((1,0)(0,-2)0) - \{(0,1)(1,2)1\}((0,0)(0,0)0)((1,0)(0,-2)0) \\
& \{(0,1)(1,2)1\}((1,0)(0,-2)0)((0,0)(0,0)0) - \{(0,1)(1,2)1\}((1,0)(0,-2)0)((0,0)(0,0)0) \\
& \{(0,1)(1,2)1\}((0,1)(1,-1)0)((0,1)(1,-1)0) - \sigma^6\{(0,1)(1,2)1\}\sigma^3((0,1)(1,-1)0)\sigma^3((0,1)(1,-1)0)
\end{aligned}$$

In order to obtain the total number of states in the **56** representation of E_6 we have to take into account that there is one state in the original theory which cancels when symmetrized. This leads again to $N_{56}=20$ as expected.

Comparing this example with the equivalent Gepner case, namely $\{2_A^2\} \ 2_A^2$, a difference arises. In [5] the appearance of the (3,0)-form coupled to the identity is used as a criterion to decide whether the Z_2 -permutation is allowed. In the case at hand the state

$$(0,0,0) \ (0,0,0) \ - \ (0,-2,-2) \ (0,-2,-2) \tag{48}$$

cancels in the symmetrized combination and thus it is not part of the spectrum. Notice that if a pair of trivial theories (i.e., $k = 0$) were added and permuted, the (3,0)-form coupled to the identity would not cancel with its symmetric form. However in the equivalent Kazama-Suzuki case, i.e. $\{(2,1)^2\} \ (2,1)^2$ the (3,0)-form is the state $(0,0)(0,-6)0$. Therefore, when coupled to the identity the symmetric combination does not cancel. This will be the case for all cosets with even m .

VI. SUMMARY AND CONCLUSIONS

We have computed the number of chiral generations in the **27** of E_6 for string models constructed with tensor products of CP_m cosets which can be described in terms of LG superpotentials. We have also extended the method developed in [4,5] to study cyclic permutation symmetries in CP_m string models which do not admit a LG formulation. This study contributes to a deeper understanding of coset theories. In particular we used explicit formulas (power expansions) for the characters of the primary fields, based on general constructions of reference [16]. In order to know the exact weights and charges of these fields just the first power of the characters is needed. Possibly this computation could be simplified by just isolating this power with some limiting process [25]. Notice that explicit expansions for the characters allow to check many of the equivalence relations among theories conjectured essentially from the chiral structures. The information needed from the original theory in order to construct the projected one, goes beyond the chiral structure. Therefore theories which are not fully conformally identical (i.e. their spectra of conformal weights and charges are not equal) may lead to different results. For example, in the case of minimal models, if equivalence at the chiral level is established then the chiral spectrum of the projected theories coincide. However the spectrum of singlets may differ [5]. An open related question remains, regarding Z_2 permutations of CP_m cosets with even m . In the example we have presented (namely $\{(2,1)^2\}(2,1)^2$), they do contribute to the massless spectrum of the orbifoldized theory, even though the equivalent Gepner case (i.e. $\{2_A^2\}2_A^2$) does not.

Moddings by phase symmetries, commuting with the cyclic permutations, may be easily incorporated into the scheme we have presented. A systematic study of cyclic permutation symmetries and phase moddings in CP_m coset models relevant for string vacua classification will be published elsewhere.

VII. ACKNOWLEDGMENTS

We are grateful to A. Font and A. Klemm for clarifying comments. This work was partially supported by Consejo Nacional de Investigaciones Científicas y Técnicas (Argentina); Universidad de Buenos Aires and Fundación Antorchas. E. A. and G. A. thank Universidad Autónoma de Madrid where part of this work was done.

VIII. APPENDIX

In this appendix we summarize some useful steps for computing the characters from [16]. There the $CP_2 = SU(3)_k / (SU(2)_{k+1} \times U(1))$ character is decomposed according to

$$\frac{SU(3)_k \times SU(2)_1 \times U(1)}{SU(2)_{k+1} \times U(1)} = \frac{SU(3)_k}{SU(2)_k \times U(1)} \times \frac{SU(2)_k \times SU(2)_1}{SU(2)_{k+1}} \times U(1)$$

Character formulae for minimal model representations of the second factor above have been built in [26]. The character for a state with charge m of the $U(1)$ theory is given by the $SU(2)_k$ theta function $\theta_{m,k}(x, y) = \sum_{n \in \mathbb{Z}} x^{k(n+m/k)^2} y^{k(n+m/k)/2}$. The character for the state $|\Lambda, \lambda = l \hat{w}_1 + \frac{w_2}{2}q, \tilde{\Lambda} = 0\rangle$ is [16]:

$$\chi_{l,q}^\Lambda(x, y) = \sum_{m=-5, \dots, 6} b_{l',m}^\Lambda(x) \chi_{l',\nu}(x) \theta_{(k+3)\frac{m-q}{6} + \frac{q}{2}; k(k+3)}(x, y)$$

where $b_{l,q}^\Lambda$ are the branching functions which can be obtained from [16] as

$$b_{l,q}^\Lambda = \prod_{n=1}^{\infty} (1-x^{-n})^{-5} \sum_{\substack{l_1, l_2 \in \mathbb{Z} \\ s_1, s_2 = 0, \dots, \infty}} (w-1)^{s_1+s_2} \epsilon(\sigma) \times x^{\frac{1}{2(k+3)} \{[\sigma(\Lambda+\rho)+(k+3)[(l_1+\frac{2}{3}s_2-\frac{1}{3}s_1)\alpha_1+(l_2+\frac{1}{3}s_2+\frac{1}{3}s_1)\alpha_2]]^2-2\}} \\ \times x^{-\frac{1}{12k} [q+k(s_1+s_2)]^2 - \frac{1}{4(k+2)} \{[l+1+(k+2)(s_2-s_1)]^2-1\}}$$

where $x = e^{2i\pi\tau}$, $\rho = \alpha_1 + \alpha_2$ and α_1, α_2 (and $\alpha_3 = \alpha_1 + \alpha_2$) denote the roots of $SU(3)$ with $\alpha_i^2 = 2$. The usual normalization $2\frac{\alpha_i \omega_j}{\alpha_i^2} = \delta_{ij}$ is taken for the weights. W is the Weyl group of $SU(3)$. Note that the sum *is not* well defined, as it depends on the order of its terms, the sums over s_i should be performed *before* the sums over l_i to get the right branching function.

Doing so, some terms cancels against others if $p_i > 0$ due to the identity [16]

$$\sum_{s=0}^{2p-1} (-1)^s x^{\frac{1}{2}(s-(p-\frac{1}{2}))^2 - \frac{1}{2}(p-\frac{1}{2})^2} = 0$$

and the sum over s_i for fixed l_i can be rewritten as

$$\sum_{s=2p}^{\infty} (-1)^s x^{\frac{1}{2}(s-(p-\frac{1}{2}))^2 - \frac{1}{2}(p-\frac{1}{2})^2} = - \sum_{s=0}^{\infty} (-1)^{-(s+1)} x^{\frac{1}{2}(-(s+1)-(p-\frac{1}{2}))^2 - \frac{1}{2}(p-\frac{1}{2})^2}$$

which amounts to the replacement $s \rightarrow -(s+1)$ and a global change of sign. The following conditions are equivalent to $p > 0$ for the s_i sum

$$\begin{aligned} s_1 : \quad & \frac{1}{3}(\alpha_2 - \alpha_1) \cdot \sigma(\Lambda + \rho) + (k+3)(l_2 - l_1) + \frac{l+1}{2} - \frac{q}{6} < 0 \\ s_2 : \quad & \frac{1}{3}(2\alpha_1 + \alpha_2) \cdot \sigma(\Lambda + \rho) + (k+3) l_1 - \frac{l+1}{2} - \frac{q}{6} < 0 \end{aligned}$$

REFERENCES

- [1] D. Gepner, Nucl. Phys. **B 296** (1987) 757; “Lectures on $N = 2$ strings”, Proceedings of the Trieste Spring School 1989, M. Green et al. (eds.), Singapore: World Scientific 1990.
- [2] Y. Kazama and H. Suzuki, Nucl. Phys. **B 321** (1989) 232.
- [3] D. Gepner, “String theory on Calabi-Yau manifolds: the 3 generation case” Princeton preprint (1987)
- [4] A. Klemm, and M. Schmidt, Phys. Lett. **B 245** (1990) 53
- [5] J. Fuchs, A. Klemm, and M. Schmidt, Ann. Phys. **214** (1992) 221
- [6] A. Font, L. E. Ibáñez, F. Quevedo and A. Sierra, Nucl. Phys. **B 337** (1990) 119.
- [7] E. Martinec, Phys. Lett. **B217** (1989) 431
- [8] C. Vafa and N. Warner, Phys. Lett. **B218** (1989) 51
- [9] W. Lerche, C. Vafa and N. Warner, Nucl. Phys. **B 324** (1989) 427
- [10] M. Kreuzer, R. Schimmrigk and H. Skarke, Nucl. Phys. **B 372** (1992) 61.
M. Kreuzer and H. Skarke, Nucl. Phys. **B 405** (1993) 305.
- [11] C. Vafa, Mod. Phys. Lett. **A 4** (1989) 1169
- [12] K. Intriligator and C. Vafa, Nucl. Phys. **B 339** (1990) 95
- [13] G. Aldazabal, I. Alekotte, A. Font and C. Núñez, Int. J. Mod. Phys. **A7** (1992) 6273.
- [14] A. Font, L. Ibáñez and F. Quevedo, Phys. Lett. **B 217** (1989) 271.
- [15] D. Gepner, Commun. Math. Phys. **142** (1991) 433.
- [16] K. Huitu, D. Nemeschansky and S. Yankielowicz, Phys. Lett. **B 246** (1990) 105
- [17] D. Gepner, Phys. Lett. **B 222** (1989) 207.

- [18] A. Sierra, “Field Identification in $N = 2$ Cosets”, preprint FTUAM/91-09 (1991).
- [19] Y. Cai and A. He, Phys. Lett. **B 252** (1990) 63.
- [20] A. N. Schellekens, Nucl. Phys. **B 366** (1991) 27.
- [21] A. N. Schellekens and S. Yankielowicz, Nucl. Phys. **B 334** (1990) 67; Int. J. Mod. Phys. **A 5** (1990) 5.
- [22] D. Gepner and Z. Qiu, Nucl. Phys. **B 285** [FS19] (1987) 423.
- [23] D. Gepner, Nucl. Phys. **B 322** (1989) 65
- [24] B.R. Greene, C. Vafa and N.P. Warner, Nucl. Phys. **B 324** (1989) 371.
J.H. Schwarz, Int. J. Mod. Phys. **A 4** (1989) 2653.
- [25] V. G. Kac, “Infinite-dimensional Lie Algebras—An Introduction”, Birkhauser, Boston, 1983, 2nd edition, Cambridge Univ. Press, Cambridge, 1985.
- [26] P. Goddard, A. Kent and D. Olive, Commun. Math. Phys. 103 (1986) 105